

SCIENTIA

Series A: *Mathematical Sciences*, Vol. ?? (2012), ??

Universidad Técnica Federico Santa María

Valparaíso, Chile

ISSN 0716-8446

© Universidad Técnica Federico Santa María 2012

## The integrals in Gradshteyn and Ryzhik. Part 23: Combination of logarithms and rational functions

Luis A. Medina and Victor H. Moll

ABSTRACT. The table of Gradshteyn and Ryzhik contains many entries where the integrand is a combination of a rational function and a logarithmic function. The proofs presented here, complete the evaluation of all entries in Section 4.231 and 4.291.

### 1. Introduction

The table of integrals [6] contains many entries of the form

$$(1.1) \quad \int_a^b R_1(x) \ln R_2(x) dx$$

where  $R_1$  and  $R_2$  are rational functions. Some of these examples have appeared in previous papers: entry **4.291.1**

$$(1.2) \quad \int_0^1 \frac{\ln(1+x)}{x} dx = \frac{\pi^2}{12}$$

as well as entry **4.291.2**

$$(1.3) \quad \int_0^1 \frac{\ln(1-x)}{x} dx = -\frac{\pi^2}{6}$$

have been established in [4], entry **4.212.7**

$$(1.4) \quad \int_1^e \frac{\ln x dx}{(1+\ln x)^2} = \frac{e}{2} - 1$$

appears in [2] and entry **4.231.11**

$$(1.5) \quad \int_0^a \frac{\ln x dx}{x^2 + a^2} = \frac{\pi \ln a}{4a} - \frac{G}{a},$$

---

2000 *Mathematics Subject Classification*. Primary 33.

*Key words and phrases*. Integrals.

The second author wishes to acknowledge the partial support of NSF-DMS 0713836.

where

$$(1.6) \quad G = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^2}$$

is the Catalan constant, has appeared in [5]. The value of entry **4.233.1**

$$(1.7) \quad \int_0^1 \frac{\ln x \, dx}{x^2 + x + 1} = \frac{2}{9} \left[ \frac{2\pi^2}{3} - \psi' \left( \frac{1}{3} \right) \right],$$

where  $\psi(x) = \Gamma'(x)/\Gamma(x)$  is the digamma function, was established in [8].

A standard trick employed in the evaluations of integrals over  $[0, \infty)$ , is to transform the interval  $[1, \infty)$  back to  $[0, 1]$  via  $t = 1/x$ . This gives

$$(1.8) \quad \int_0^{\infty} R(x) \ln x \, dx = \int_0^1 \left[ R(x) - \frac{1}{x^2} R \left( \frac{1}{x} \right) \right] dx.$$

In particular, if the rational function satisfies

$$(1.9) \quad R \left( \frac{1}{x} \right) = x^2 R(x),$$

then

$$(1.10) \quad \int_0^{\infty} R(x) \ln x \, dx = 0.$$

This is the case for  $R(x) = \frac{1+x^2}{(1-x^2)^2}$  and (1.10) appears as entry **4.234.3** in [6].

The goal of this paper is to present a systematic evaluation of the entries in [6] of the form (1.1).

## 2. Combinations of logarithms and linear rational functions

EXAMPLE 2.1. Entry **4.291.3** states that

$$(2.1) \quad \int_0^{1/2} \frac{\ln(1-x)}{x} \, dx = \frac{\ln^2 2}{2} - \frac{\pi^2}{12}.$$

To evaluate this integral let  $t = -\ln(1-x)$  to produce

$$(2.2) \quad \int_0^{1/2} \frac{\ln(1-x)}{x} \, dx = - \int_0^{\ln 2} \frac{te^{-t} \, dt}{1 - e^{-t}}.$$

This last integral can be written as

$$(2.3) \quad \int_0^{\ln 2} t \, dt - \int_0^{\ln 2} \frac{t \, dt}{1 - e^{-t}}.$$

The first integral is elementary and has value  $\frac{1}{2} \ln^2 2$ . The second integral was evaluated as  $\pi^2/12$  in [3].

EXAMPLE 2.2. The change of variables  $t = x/2$  converts (2.1) to

$$(2.4) \quad \int_0^{1/2} \ln\left(1 - \frac{t}{2}\right) \frac{dt}{t} = \frac{\ln^2 2}{2} - \frac{\pi^2}{12}.$$

This is entry **4.291.4** of [6].

EXAMPLE 2.3. Entry **4.291.5** states that

$$(2.5) \quad \int_0^1 \ln\left(\frac{1+x}{2}\right) \frac{dx}{1-x} = \frac{\ln^2 2}{2} - \frac{\pi^2}{12}.$$

To evaluate this entry, let  $u = (1-x)/2$  to reduce it to (2.1)

EXAMPLE 2.4. Differentiating

$$(2.6) \quad \int_0^1 (1+x)^{-a} dx = \frac{2^{-a}(2^a - 2)}{a-1}$$

with respect to  $a$  gives

$$(2.7) \quad \int_0^1 (1+x)^{-a} \ln(1+x) dx = \frac{1}{(a-1)^2} (2^{-a}(-2 + 2^a + 2 \ln 2 - 2a \ln 2)).$$

Now let  $a \rightarrow 1$  to obtain

$$(2.8) \quad \int_0^1 \frac{\ln(1+x)}{1+x} dx = \frac{1}{2} \ln^2 2.$$

This is entry **4.291.6**.

EXAMPLE 2.5. The partial fraction decomposition

$$(2.9) \quad \frac{1}{x(1+x)} = \frac{1}{x} - \frac{1}{1+x}$$

gives

$$(2.10) \quad \int_0^1 \frac{\ln(1+x)}{x(1+x)} dx = \int_0^1 \frac{\ln(1+x)}{x} dx - \int_0^1 \frac{\ln(1+x)}{1+x} dx.$$

The first integral is entry **4.291.1** and it has value  $\pi^2/12$  as shown in [4]. The second integral is  $\frac{1}{2} \ln^2 2$  as established in Example 2.4. This gives entry **4.291.12**

$$(2.11) \quad \int_0^1 \frac{\ln(1+x)}{x(1+x)} dx = \frac{\pi^2}{12} - \frac{1}{2} \ln^2 2.$$

EXAMPLE 2.6. Entry **4.291.13** is

$$(2.12) \quad \int_0^\infty \frac{\ln(1+x) dx}{x(1+x)} = \frac{\pi^2}{6}.$$

Split the integral over  $[0, 1]$  and  $[1, \infty)$  and make the change of variables  $t = 1/x$  in the second part. This gives

$$(2.13) \quad \int_0^\infty \frac{\ln(1+x) dx}{x(1+x)} = \int_0^1 \frac{\ln(1+x) dx}{x(1+x)} + \int_0^1 \frac{\ln(1+t) - \ln t}{1+t} dt.$$

Expand the first integral in partial fractions to obtain

$$(2.14) \quad \int_0^\infty \frac{\ln(1+x) dx}{x(1+x)} = \int_0^1 \frac{\ln(1+x)}{x} dx - \int_0^1 \frac{\ln x}{1+x} dx.$$

Integrate by parts the second integral to obtain

$$(2.15) \quad \int_0^\infty \frac{\ln(1+x) dx}{x(1+x)} = 2 \int_0^1 \frac{\ln(1+x)}{x} dx.$$

The evaluation

$$(2.16) \quad \int_0^1 \frac{\ln(1+x)}{x} dx = \frac{\pi^2}{12}$$

that appears as entry **4.291.1** has been established in [4].

### 3. Combinations of logarithms and rational functions with denominators that are squares of linear terms

This section evaluates integrals of the form

$$(3.1) \quad \int_a^b R_2(x) \ln R_1(x) dx$$

where  $R_1, R_2$  are rational functions and the denominator of  $R_2$  is a quadratic polynomial of the form  $(cx+d)^2$ .

EXAMPLE 3.1. Entry **4.291.14** is

$$(3.2) \quad \int_0^1 \frac{\ln(1+x)}{(ax+b)^2} dx = \frac{1}{a(a-b)} \ln \frac{a+b}{b} + \frac{2 \ln 2}{b^2 - a^2}$$

and

$$(3.3) \quad \int_0^1 \frac{\ln(1+x) dx}{(x+1)^2} = \frac{1 - \ln 2}{2}$$

gives the value when  $a = b$ , after scaling.

To evaluate the first case, integrate by parts to get

$$(3.4) \quad \int_0^1 \frac{\ln(1+x)}{(ax+b)^2} dx = -\frac{\ln 2}{a(a+b)} + \frac{1}{a} \int_0^1 \frac{dx}{(1+x)(ax+b)}.$$

The result now follows by expanding the second integrand in partial fractions.

The case  $a = b$  is obtained by a direct integration by parts:

$$(3.5) \quad \int_0^1 \frac{\ln(1+x)}{(1+x)^2} dx = -\frac{\ln 2}{2} + \int_0^1 \frac{dx}{(1+x)^2}.$$

This last integral is  $1/2$  and the result has been established.

The same procedure gives entry **4.291.20**:

$$(3.6) \quad \int_0^1 \frac{\ln(ax+b)}{(1+x)^2} dx = \frac{1}{2(a-b)} [(a+b) \ln(a+b) - 2b \ln b - 2a \ln 2],$$

for  $a \neq b$ .

EXAMPLE 3.2. The partial fraction decomposition

$$(3.7) \quad \frac{1-x^2}{(ax+b)^2(bx+a)^2} = \frac{1}{a^2-b^2} \left[ \frac{1}{(ax+b)^2} - \frac{1}{(bx+a)^2} \right]$$

and Example 3.1 gives the evaluation of entry **4.291.25**:

$$\int_0^1 \frac{(1-x^2)\ln(1+x)dx}{(ax+b)^2(bx+a)^2} = \frac{1}{(a^2-b^2)(a-b)} \left[ \frac{a+b}{ab} \ln(a+b) - \frac{\ln b}{a} - \frac{\ln a}{b} \right] - \frac{4\ln 2}{(a^2-b^2)^2}.$$

The answer may be written in the more compact form

$$(3.8) \quad \frac{-a^2 \ln a - b[b \ln b + a \ln(16ab)] + (a+b)^2 \ln(a+b)}{ab(a^2-b^2)^2},$$

but this form hides the symmetry of the integral.

EXAMPLE 3.3. Entry **4.291.15** is

$$(3.9) \quad \int_0^\infty \frac{\ln(1+x)dx}{(ax+b)^2} = \frac{\ln a - \ln b}{a(a-b)}$$

for  $a \neq b$ . In the case  $a = b$ , the integral scales to

$$(3.10) \quad \int_0^\infty \frac{\ln(1+x)dx}{(1+x)^2} = 1.$$

To evaluate this entry, integrate by parts to obtain

$$(3.11) \quad \int_0^\infty \frac{\ln(1+x)dx}{(ax+b)^2} = \frac{1}{a} \int_0^\infty \frac{dx}{(1+x)(ax+b)}.$$

This last integral is evaluated by using the partial fraction decomposition

$$(3.12) \quad \frac{1}{(1+x)(ax+b)} = \frac{1}{b-a} \left( \frac{1}{1+x} - \frac{a}{ax+b} \right).$$

Integration by parts in the case  $a = b$  (taken to be 1 by scaling) gives

$$(3.13) \quad \int_0^\infty \frac{\ln(1+x)dx}{(1+x)^2} = \int_0^\infty \frac{dx}{(1+x)^2} = 1.$$

The same procedure gives entry **4.291.21**:

$$(3.14) \quad \int_0^\infty \frac{\ln(ax+b)dx}{(1+x)^2} = \frac{a \ln a - b \ln b}{a-b}.$$

for  $a \neq b$ . The value of entry **4.291.17**:

$$(3.15) \quad \int_0^\infty \frac{\ln(a+x)}{(b+x)^2} dx = \frac{a \ln a - b \ln b}{b(a-b)}$$

is obtained from (3.14) by the change of variables  $x = bt$ .

EXAMPLE 3.4. The partial fraction decomposition (3.7) given in Example 3.2 produces the value of entry **4.291.26**

$$(3.16) \quad \int_0^\infty \frac{(1-x^2)\ln(1+x)dx}{(ax+b)^2(bx+a)^2} = \frac{\ln b - \ln a}{ab(a^2 - b^2)}$$

form Example 3.3.

#### 4. Combinations of logarithms and rational functions with quadratic denominators

This section considers integrals of the form (1.1) where the denominator of  $R_2(x)$  is a polynomial of degree 2 with non-real roots.

EXAMPLE 4.1. Entry **4.291.8** states that

$$(4.1) \quad \int_0^1 \frac{\ln(1+x)dx}{1+x^2} = \frac{\pi}{8} \ln 2.$$

The proof of this evaluation is based on some entries of [6] that have been established in [4]. The reader is invited to provide a direct proof.

The change of variables  $x = \tan \varphi$  gives

$$\begin{aligned} \int_0^1 \frac{\ln(1+x)dx}{1+x^2} &= \int_0^{\pi/4} \ln(1+\tan \varphi)d\varphi \\ &= \int_0^{\pi/4} \ln(\sin \varphi + \cos \varphi)d\varphi - \int_0^{\pi/4} \ln \cos \varphi d\varphi. \end{aligned}$$

The value

$$\int_0^{\pi/4} \ln(\sin \varphi + \cos \varphi)d\varphi = -\frac{\pi}{8} \ln 2 + \frac{G}{2}$$

is entry **4.225.2** and

$$\int_0^{\pi/4} \ln \cos \varphi d\varphi = -\frac{\pi}{4} \ln 2 + \frac{G}{2}$$

is entry **4.224.5**. Both examples are evaluated in [4]. This gives the result.

The same technique gives entry **4.291.10**

$$(4.2) \quad \int_0^1 \frac{\ln(1-x)dx}{1+x^2} = \frac{\pi}{8} \ln 2 - G.$$

This time, entry **4.225.1**

$$\int_0^{\pi/4} \ln(\cos \varphi - \sin \varphi)d\varphi = -\frac{\pi}{8} \ln 2 - \frac{G}{2}$$

is employed.

EXAMPLE 4.2. Entry **4.291.9**

$$(4.3) \quad \int_0^\infty \frac{\ln(1+x)dx}{1+x^2} = \frac{\pi}{4} \ln 2 + G$$

is equivalent, via  $x = \tan \varphi$ , to the identity

$$(4.4) \quad \int_0^{\pi/2} \ln(\sin \varphi + \cos \varphi) d\varphi - \int_0^{\pi/2} \ln \cos \varphi d\varphi = \frac{\pi}{4} \ln 2 + G.$$

The first integral is entry **4.225.2** and it has the value  $-\frac{1}{4}\pi \ln 2 + G$ ; the second integral is entry **4.224.6** with value  $-\frac{1}{2}\pi \ln 2$ . Both of these examples have been established in [4].

EXAMPLE 4.3. The change of variables  $t = 1/x$  gives

$$(4.5) \quad \int_1^{\infty} \frac{\ln(x-1) dx}{1+x^2} = \int_0^1 \frac{\ln(1-t) dt}{1+t^2} - \int_0^1 \frac{\ln t dt}{1+t^2}.$$

The first integral has the value  $\frac{1}{8}\pi \ln 2 - G$  and it appears as entry **4.291.10** (it has been established as (4.2)). The second integral is the special case  $a = 1$  of (1.5). This gives the value of entry **4.291.11**:

$$(4.6) \quad \int_1^{\infty} \frac{\ln(x-1) dx}{1+x^2} = \frac{\pi}{8} \ln 2.$$

EXAMPLE 4.4. A small number of entries in [6] can be evaluated from entry **4.231.9**

$$(4.7) \quad \int_0^{\infty} \frac{\ln x dx}{x^2 + q^2} = \frac{\pi \ln q}{2q},$$

evaluated in [4]. Expanding in partial fractions gives the identity

$$(4.8) \quad \int_0^{\infty} \frac{\ln x dx}{(x^2 + a^2)(x^2 + b^2)} = \frac{\pi}{2(b^2 - a^2)} \left( \frac{\ln a}{a} - \frac{\ln b}{b} \right).$$

This provides the evaluation of entry **4.234.6**

$$(4.9) \quad \int_0^{\infty} \frac{\ln x dx}{(a^2 + b^2 x^2)(1 + x^2)} = \frac{\pi b}{2a(b^2 - a^2)} \ln \frac{a}{b}$$

via the relation

$$(4.10) \quad \int_0^{\infty} \frac{\ln x dx}{(a^2 + b^2 x^2)(1 + x^2)} = \frac{1}{b^2} \int_0^{\infty} \frac{\ln x dx}{(x^2 + a^2/b^2)(x^2 + 1)},$$

entry **4.234.7**

$$(4.11) \quad \int_0^{\infty} \frac{\ln x dx}{(x^2 + a^2)(1 + b^2 x^2)} = \frac{\pi}{2(1 - a^2 b^2)} \left( \frac{\ln a}{a} + b \ln b \right)$$

via the relation

$$(4.12) \quad \int_0^{\infty} \frac{\ln x dx}{(x^2 + a^2)(1 + b^2 x^2)} = \frac{1}{b^2} \int_0^{\infty} \frac{\ln x dx}{(x^2 + a^2)(x^2 + 1/b^2)},$$

and finally, entry **4.234.8**

$$(4.13) \quad \int_0^{\infty} \frac{x^2 \ln x dx}{(a^2 + b^2 x^2)(1 + x^2)} = \frac{\pi a}{2b(b^2 - a^2)} \ln \frac{b}{a}$$

using the partial fraction decomposition

$$(4.14) \quad \frac{x^2}{(a^2 + b^2x^2)(1+x^2)} = \frac{1}{(b^2 - a^2)} \frac{1}{x^2 + 1} - \frac{a^2}{b^2(b^2 - a^2)} \frac{1}{x^2 + a^2/b^2}.$$

The details are left to the reader.

### 5. An example via recurrences

The integral

$$(5.1) \quad F_n(s) = \int_0^1 x^n(1+x)^s dx$$

for  $n \in \mathbb{N}$  and  $s \in \mathbb{R}$ , is integrated by parts (with  $u = x^n(1+x)$  and  $dv = (1+x)^{s-1} dx$ ), to produce the recurrence

$$(5.2) \quad F_n(s) = \frac{2^{s+1}}{n+s+1} - \frac{n}{n+s+1} F_{n-1}(s).$$

The initial condition is

$$(5.3) \quad F_0(s) = \int_0^1 (1+x)^s dx = \frac{2^{s+1} - 1}{s+1}.$$

The recurrence permits the evaluation of  $F_n(s)$ , for any fixed  $n \in \mathbb{N}$ . For instance,

$$\begin{aligned} F_1(s) &= \frac{s2^{s+1} + 1}{(s+1)(s+2)} \\ F_2(s) &= \frac{2 [2^s(s^2 + s + 2) - 1]}{(s+1)(s+2)(s+3)} \\ F_3(s) &= \frac{2 [2^s(s^3 + 3s^2 + 8s) + 3]}{(s+1)(s+2)(s+3)(s+4)}. \end{aligned}$$

Differentiating (5.2) produces a recurrence for

$$(5.4) \quad G_n(s) = \int_0^1 \frac{x^n \ln(1+x)}{(1+x)^s} dx$$

in the form

$$(5.5) \quad G_n(s) = -\frac{2^{1-s}}{(n+1-s)^2} + \frac{2^{1-s} \ln 2}{n+1-s} + \frac{n}{(n-s+1)^2} F_n(-s) - \frac{n}{n-s+1} G_{n-1}(s).$$

This produces the value of  $G_n(s)$ , starting from

$$(5.6) \quad G_0(s) = \int_0^1 \frac{\ln(1+x)}{(1+x)^s} dx = \frac{2^{1-s} \ln 2}{1-s} - \frac{2^{1-s} - 1}{(1-s)^2}.$$

For example,

$$(5.7) \quad G_1(s) = \frac{2^s(2s-3) - 2 \ln 2s^3 + 2(3 \ln 2 - 1)s^2 - 4 \ln 2s + 4}{2^s(s-1)^2(s-2)^2}.$$



EXAMPLE 5.1. Entry **4.291.23** in [6] states that

$$(5.8) \quad \int_0^1 \ln(1+x) \frac{1+x^2}{(1+x)^4} dx = -\frac{\ln 2}{3} + \frac{23}{72}.$$

This corresponds to the value  $G_0(4) + G_2(4)$ . The recurrence (5.5) gives the required data to verify this entry.

### 6. An elementary example

Integrals of the form

$$(6.1) \quad \int_a^b \ln R_1(x) \frac{d}{dx} R_2(x) dx$$

for rational functions  $R_1, R_2$  can be reduced to the integration of a rational function. Indeed, integration by parts yields

$$(6.2) \quad \int_a^b \ln R_1(x) \frac{d}{dx} R_2(x) dx = \text{boundary terms} - \int_a^b R_3(x) dx$$

with  $R_3 = R_1' R_2 / R_1$ .

EXAMPLE 6.1. Entry **4.291.27** states that

$$(6.3) \quad \int_0^1 \ln(1+ax) \frac{1-x^2}{(1+x^2)^2} dx = \frac{(1+a)^2 \ln(1+a)}{1+a^2} - \frac{\ln 2}{2} - \frac{a}{1+a^2} - \frac{\pi}{4} \frac{a^2}{1+a^2}.$$

This example fits the pattern described above, since

$$(6.4) \quad \frac{1-x^2}{(1+x^2)^2} = \frac{d}{dx} \frac{x}{1+x^2}.$$

Therefore

$$\begin{aligned} \int_0^1 \ln(1+ax) \frac{1-x^2}{(1+x^2)^2} dx &= \int_0^1 \ln(1+ax) \frac{d}{dx} \frac{x}{1+x^2} dx \\ &= \frac{\ln(1+a)}{2} - a \int_0^1 \frac{x dx}{(1+x^2)(1+ax)}. \end{aligned}$$

The partial fraction decomposition

$$\frac{x}{(1+x^2)(1+ax)} = -\frac{a}{1+a^2} \frac{1}{1+ax} + \frac{a}{1+a^2} \frac{1}{1+x^2} + \frac{1}{1+a^2} \frac{x}{1+x^2}$$

and the evaluation of the remaining elementary integrals completes the solution to this problem.

EXAMPLE 6.2. Entry **4.291.28**

$$(6.5) \quad \int_0^\infty \ln(a+x) \frac{b^2-x^2}{(b^2+x^2)^2} dx = \frac{1}{a^2+b^2} \left( a \ln \frac{b}{a} - \frac{\pi b}{2} \right)$$

also fits the pattern in this section since

$$(6.6) \quad \frac{d}{dx} \frac{x}{x^2+b^2} = \frac{b^2-x^2}{(b^2+x^2)^2}.$$

Integrating by parts and checking that the boundary terms vanish, produces

$$(6.7) \quad \int_0^\infty \ln(a+x) \frac{b^2-x^2}{(b^2+x^2)^2} dx = - \int_0^\infty \frac{x dx}{(x^2+b^2)(x+a)}.$$

It is convenient to introduce the scaling  $x = bt$  to transform the last integral to

$$(6.8) \quad \int_0^\infty \frac{x dx}{(x^2+b^2)(x+a)} = \frac{1}{b} \int_0^\infty \frac{t dt}{(1+t^2)(t+c)}$$

with  $c = a/b$ . The evaluation is completed using the partial fraction decomposition

$$\frac{t}{(t^2+1)(t+c)} = -\frac{c}{c^2+1} \frac{1}{t+c} + \frac{1}{1+c^2} \frac{1}{t^2+1} + \frac{c}{c^2+1} \frac{t}{t^2+1}$$

and integrating from  $t = 0$  to  $t = N$  and taking the limit as  $N \rightarrow \infty$ . The reader will easily check that the divergent pieces, coming from  $1/(t+c)$  and  $t/(t^2+1)$  cancel out.

EXAMPLE 6.3. Entry **4.291.29** appears as

$$(6.9) \quad \int_0^\infty \ln^2(a-x) \frac{b^2-x^2}{(b^2+x^2)^2} dx = \frac{2}{a^2+b^2} \left( a \ln \frac{a}{b} - \frac{\pi b}{2} \right)$$

but it should be written as

$$(6.10) \quad \int_0^\infty \ln [(a-x)^2] \frac{b^2-x^2}{(b^2+x^2)^2} dx = \frac{2}{a^2+b^2} \left( a \ln \frac{a}{b} - \frac{\pi b}{2} \right).$$

This is a singular integral and the value should be interpreted as a Cauchy principal value

$$\begin{aligned} \int_0^\infty \ln [(a-x)^2] \frac{b^2-x^2}{(b^2+x^2)^2} dx &= \\ \lim_{\varepsilon \rightarrow 0} \int_0^{a-\varepsilon} \ln [(a-x)^2] \frac{b^2-x^2}{(b^2+x^2)^2} dx &+ \int_{a+\varepsilon}^\infty \ln [(a-x)^2] \frac{b^2-x^2}{(b^2+x^2)^2} dx. \end{aligned}$$

The first integral is

$$\begin{aligned} \int_0^{a-\varepsilon} \ln [(a-x)^2] \frac{b^2-x^2}{(b^2+x^2)^2} dx &= \int_0^{a-\varepsilon} 2 \ln(a-x) \frac{d}{dx} \frac{x}{x^2+b^2} dx \\ &= \frac{2(a-\varepsilon)}{(a-\varepsilon)^2+b^2} \ln \varepsilon + \int_0^{a-\varepsilon} \frac{2x dx}{(a-x)(x^2+b^2)}, \end{aligned}$$

after integration by parts. The second integral produces

$$\begin{aligned} \int_{a+\varepsilon}^\infty \ln [(a-x)^2] \frac{b^2-x^2}{(b^2+x^2)^2} dx &= \int_{a+\varepsilon}^\infty 2 \ln(x-a) \frac{d}{dx} \frac{x}{x^2+b^2} dx \\ &= -\frac{2(a+\varepsilon)}{(a+\varepsilon)^2+b^2} \ln \varepsilon + \int_{a+\varepsilon}^\infty \frac{2x dx}{(x-a)(x^2+b^2)}. \end{aligned}$$

The reader will check that the boundary terms vanish as  $\varepsilon \rightarrow 0$ . This produces

$$(6.11) \quad \int_0^\infty \ln[(a-x)^2] \frac{b^2-x^2}{(b^2+x^2)^2} dx = \lim_{\varepsilon \rightarrow 0} \int_0^{a-\varepsilon} \frac{2x dx}{(a-x)(x^2+b^2)} + \int_{a+\varepsilon}^\infty \frac{2x dx}{(a-x)(x^2+b^2)}.$$

The partial fraction decomposition

$$(6.12) \quad \frac{2x}{(a-x)(x^2+b^2)} = -\frac{2a}{a^2+b^2} \frac{1}{x-a} - \frac{2b}{a^2+b^2} \frac{b}{x^2+b^2} + \frac{a}{a^2+b^2} \frac{2x}{x^2+b^2}$$

gives

$$\int_0^{a-\varepsilon} \frac{2x dx}{(a-x)(x^2+b^2)} = \frac{2a}{a^2+b^2} [\ln a - \ln \varepsilon] - \frac{2b}{a^2+b^2} \tan^{-1} \frac{a-\varepsilon}{b} + \frac{a}{a^2+b^2} [\ln[(a-\varepsilon)^2+b^2] - 2 \ln b].$$

A similar computation yields

$$\int_{a+\varepsilon}^N \frac{2x dx}{(a-x)(x^2+b^2)} = \frac{a}{a^2+b^2} \{ \ln(N^2+b^2) - 2 \ln(N-a) + 2 \ln \varepsilon - \ln[(a+\varepsilon)^2+b^2] \} + \frac{2b}{a^2+b^2} \left[ \tan^{-1} \left( \frac{a+\varepsilon}{b} \right) - \tan^{-1} \left( \frac{N}{b} \right) \right].$$

Now let  $N \rightarrow \infty$  and use  $\ln(N^2+b^2) - 2 \ln(N-a) \rightarrow 0$  to obtain

$$\int_{a+\varepsilon}^\infty \frac{2x dx}{(a-x)(x^2+b^2)} = \frac{a}{a^2+b^2} \{ 2 \ln \varepsilon - \ln[(a+\varepsilon)^2+b^2] \} + \frac{2b}{a^2+b^2} \left[ \tan^{-1} \left( \frac{a+\varepsilon}{b} \right) - \frac{\pi}{2} \right].$$

Observe that the singular terms in (6.11), namely those containing the factor  $\ln \varepsilon$ , cancel out. The remaining terms produce the stated answer as  $\varepsilon \rightarrow 0$ . This completes the evaluation.

EXAMPLE 6.4. Entry **4.291.30** written as

$$(6.13) \quad \int_0^\infty \ln[(a-x)^2] \frac{x dx}{(b^2+x^2)^2} = \frac{1}{a^2+b^2} \left( \ln b - \frac{\pi a}{2b} + \frac{a^2}{b^2} \ln a \right)$$

is evaluated as Example 6.3. Start with the identity

$$(6.14) \quad \frac{d}{dx} \left( -\frac{1}{2(x^2+b^2)} \right) = \frac{x}{(x^2+b^2)^2}$$

and then proceed as before. The details are elementary and they are left to the reader.

### 7. Some parametric examples

This section considers some entries of [6] that depend on a parameter.

EXAMPLE 7.1. Entry **4.291.18** states that

$$(7.1) \quad \int_0^a \frac{\ln(1+ax) dx}{1+x^2} = \frac{1}{2} \tan^{-1} a \ln(1+a^2).$$

Differentiating the left-hand side with respect to  $a$  gives

$$(7.2) \quad \frac{\ln(1+a^2)}{1+a^2} + \int_0^a \frac{x dx}{(1+ax)(1+x^2)}.$$

The verification of this entry will start with the evaluation of the rational integral

$$(7.3) \quad R(a) := \int_0^a \frac{x dx}{(1+ax)(1+x^2)}.$$

The partial fraction decomposition

$$(7.4) \quad \frac{x}{(1+ax)(1+x^2)} = -\frac{1}{1+a^2} \frac{a}{1+ax} + \frac{a}{1+a^2} \frac{1}{1+x^2} + \frac{1}{2(1+a^2)} \frac{2x}{1+x^2}$$

gives

$$(7.5) \quad R(a) = -\frac{\ln(1+a^2)}{1+a^2} + \frac{a}{1+a^2} \tan^{-1} a + \frac{\ln(1+a^2)}{2(1+a^2)}.$$

Motivated by the expression in the entry being evaluated, observe that

$$(7.6) \quad \int_0^a \frac{x dx}{(1+ax)(1+x^2)} + \frac{\ln(1+a^2)}{1+a^2} = \frac{1}{2} \frac{d}{da} [\tan^{-1} a \ln(1+a^2)].$$

Now integrate this identity from 0 to  $a$  to obtain

$$(7.7) \quad \int_0^a \left[ \int_0^b \frac{x dx}{(1+bx)(1+x^2)} + \frac{\ln(1+b^2)}{1+b^2} \right] db + \int_0^a \frac{\ln(1+b^2)}{1+b^2} db = \frac{1}{2} \tan^{-1} a \ln(1+a^2).$$

Exchange the order of integration to produce

$$\begin{aligned} \int_0^a \int_0^b \frac{x dx}{(1+bx)(1+x^2)} db &= \int_0^a \frac{x}{1+x^2} \int_x^a \frac{db}{1+bx} dx \\ &= \int_0^a \frac{1}{1+x^2} [\ln(1+ax) - \ln(1+x^2)] dx. \end{aligned}$$

The result now follows from (7.7).

EXAMPLE 7.2. Entry **4.291.16** states that

$$(7.8) \quad \int_0^1 \frac{\ln(a+x) dx}{a+x^2} = \frac{1}{2\sqrt{a}} \cot^{-1} \sqrt{a} \ln[a(1+a)].$$

The change of variables  $x = \sqrt{at}$  gives

$$(7.9) \quad \int_0^1 \frac{\ln(a+x) dx}{a+x^2} = \frac{1}{\sqrt{a}} \left[ \ln a \int_0^{1/\sqrt{a}} \frac{dt}{1+t^2} + \int_0^{1/\sqrt{a}} \frac{\ln(1+t/\sqrt{a})}{1+t^2} dt \right].$$

The first integral is elementary and the second one corresponds to (7.1).

EXAMPLE 7.3. Entry **4.291.19** states that

$$(7.10) \quad \int_0^1 \frac{\ln(1+ax) dx}{1+ax^2} = \frac{1}{2\sqrt{a}} \tan^{-1} \sqrt{a} \ln(1+a).$$

This follows directly from (7.1) by the change of variables  $x = t/\sqrt{a}$  and replacing  $a$  by  $\sqrt{a}$ .

EXAMPLE 7.4. Entry **4.291.7** is the identity

$$(7.11) \quad \int_0^\infty \frac{\ln(1+ax) dx}{1+x^2} = \frac{\pi}{4} \ln(1+a^2) - \int_0^a \frac{\ln u du}{1+u^2}.$$

Differentiating the left-hand side gives

$$\begin{aligned} \frac{d}{da} \int_0^\infty \frac{\ln(1+ax) dx}{1+x^2} &= \int_0^\infty \frac{x dx}{(1+ax)(1+x^2)} \\ &= \frac{\pi}{2} \frac{a}{1+a^2} - \frac{\ln a}{1+a^2}, \end{aligned}$$

where the last evaluation is established by partial fractions. The result now follows by integrating back with respect to  $a$ .

REMARK 7.1. The current version of **Mathematica** gives

$$\int_0^a \frac{\ln x dx}{1+x^2} = \tan^{-1} a \ln a - \frac{i}{2} \text{PolyLog}[2, -ia] + \frac{i}{2} \text{PolyLog}[2, ia]$$

but is unable to provide an analytic expression for the integral

$$\int_0^\infty \frac{\ln(1+ax) dx}{1+x^2}.$$

Entries of [6] that can be evaluated in terms of polylogarithms will be described in a future publication.

EXAMPLE 7.5. Entry **4.291.24** states that

$$\int_0^1 \frac{(1+x^2) \ln(1+x)}{(a^2+x^2)(1+a^2x^2)} dx = \frac{1}{2a(1+a^2)} \left[ \frac{\pi}{2} \ln(1+a^2) - 2 \tan^{-1} a \ln a \right].$$

The evaluation of this entry starts with the partial fraction decomposition

$$(7.12) \quad \frac{1+x^2}{(a^2+x^2)(1+a^2x^2)} = \frac{1}{1+a^2} \left[ \frac{1}{x^2+a^2} + \frac{1}{1+a^2x^2} \right]$$

that yields the identity

$$\int_0^1 \frac{(1+x^2) \ln(1+x)}{(a^2+x^2)(1+a^2x^2)} dx = \frac{1}{1+a^2} \left[ \int_0^1 \frac{\ln(1+x) dx}{x^2+a^2} + \int_0^1 \frac{\ln(1+x) dx}{1+a^2x^2} \right],$$

and the change of variables  $t = 1/x$  then produces

$$\int_0^1 \frac{\ln(1+x) dx}{1+a^2x^2} = \int_1^\infty \frac{\ln(1+t) dt}{t^2+a^2} - \int_1^\infty \frac{\ln t dt}{t^2+a^2}.$$

Therefore

$$\int_0^1 \frac{(1+x^2) \ln(1+x)}{(a^2+x^2)(1+a^2x^2)} dx = \frac{1}{1+a^2} \left[ \int_0^\infty \frac{\ln(1+x) dx}{x^2+a^2} - \int_1^\infty \frac{\ln x dx}{x^2+a^2} \right].$$

The change of variables  $x = at$  and Example 7.4 give

$$\begin{aligned} \int_0^\infty \frac{\ln(1+x) dx}{x^2+a^2} &= \frac{1}{a} \int_0^\infty \frac{\ln(1+at) dt}{1+t^2} \\ &= \frac{\pi}{4a} \ln(1+a^2) - \frac{1}{a} \int_0^a \frac{\ln t dt}{1+t^2}. \end{aligned}$$

Therefore

$$(7.13) \quad \int_0^1 \frac{(1+x^2) \ln(1+x)}{(a^2+x^2)(1+a^2x^2)} dx = \frac{1}{1+a^2} \left[ \frac{\pi}{4a} \ln(1+a^2) - \frac{1}{a} \int_0^a \frac{\ln x dx}{1+x^2} - \int_1^\infty \frac{\ln x dx}{x^2+a^2} \right].$$

The change of variables  $x = at$  gives

$$\begin{aligned} \int_1^\infty \frac{\ln x dx}{x^2+a^2} &= \frac{\ln a}{a} \int_{1/a}^\infty \frac{dt}{1+t^2} + \frac{1}{a} \int_{1/a}^\infty \frac{\ln t dt}{1+t^2} \\ &= \frac{\ln a}{a} \int_{1/a}^\infty \frac{dt}{1+t^2} - \frac{1}{a} \int_0^a \frac{\ln u du}{1+u^2}, \end{aligned}$$

after the change of variables  $u = 1/t$  in the last integral. Replacing in (7.13) gives the result.

EXAMPLE 7.6. The last entry of [6] discussed here is **4.291.22**

$$\int_0^\infty \frac{x \ln(a+x)}{(b^2+x^2)^2} dx = \frac{1}{2(a^2+b^2)} \left( \ln b + \frac{\pi a}{2b} + \frac{a^2}{b^2} \ln a \right).$$

As before, start with the identity

$$(7.14) \quad \frac{x}{(x^2+b^2)^2} = -\frac{d}{dx} \frac{1}{2(x^2+b^2)}$$

and integrate by parts to produce

$$\int_0^\infty \frac{x \ln(a+x)}{(b^2+x^2)^2} dx = \frac{\ln a}{2b^2} + \frac{1}{2} \int_0^\infty \frac{dx}{(x+a)(x^2+b^2)}.$$

This last integral is evaluated by the method of partial fractions to obtain the result.

**Summary.** The examples presented here, complete the evaluation of every entry in Section 4.291 of the table [6]. The entries not appearing here have been presented in [4, 5, 7].

### 8. Integrals yielding partial sums of the zeta function

Some entries of [6] contain as the integrand the product of  $\ln x$  and a rational function coming from manipulations of a geometric series. This section presents the evaluation of some of these examples. These evaluations can be written in terms of the Riemann zeta function

$$(8.1) \quad \zeta(s) = \sum_{k=1}^{\infty} \frac{1}{n^s}$$

and the generalized harmonic numbers

$$(8.2) \quad H_{n,m} = \sum_{k=1}^n \frac{1}{k^m}.$$

EXAMPLE 8.1. Entry **4.231.18** states that

$$(8.3) \quad \int_0^1 \frac{1-x^{n+1}}{(1-x)^2} \ln x \, dx = -\frac{(n+1)\pi^2}{6} + \sum_{k=1}^n \frac{n-k+1}{k^2}.$$

This can be expressed as

$$(8.4) \quad \int_0^1 \frac{1-x^{n+1}}{(1-x)^2} \ln x \, dx = -(n+1)\zeta(2) + (n+1)H_{n,2} - H_{n,1}.$$

The evaluation begins with the identity

$$(8.5) \quad \frac{1}{(1-x)^2} = \sum_{k=0}^{\infty} (k+1)x^k$$

and its shift

$$(8.6) \quad \frac{1-x^{n+1}}{(1-x)^2} = \sum_{k=0}^n (k+1)x^k + (n+1) \sum_{k=n+1}^{\infty} x^k.$$

Integrate term by term and use the value

$$(8.7) \quad \int_0^1 x^k \ln x \, dx = -\frac{1}{(k+1)^2}$$

to obtain

$$(8.8) \quad \int_0^1 \frac{1-x^{n+1}}{(1-x)^2} \ln x \, dx = -\sum_{k=0}^n \frac{1}{k+1} - (n+1) \sum_{k=n+1}^{\infty} \frac{1}{(k+1)^2}.$$

This can now be transformed to the form stated in [6].

EXAMPLE 8.2. Entry **4.262.7**

$$(8.9) \quad \int_0^1 \frac{1-x^{n+1}}{(1-x)^2} (\ln x)^3 \, dx = -\frac{(n+1)\pi^4}{15} + 6 \sum_{k=1}^n \frac{n-k+1}{k^4}$$

is obtained by using (8.6), the identity

$$(8.10) \quad \int_0^1 (\ln x)^3 x^k \, dx = -\frac{6}{(k+1)^4},$$

and the value

$$(8.11) \quad \sum_{k=1}^{\infty} \frac{1}{k^4} = \zeta(4) = \frac{\pi^4}{90}.$$

EXAMPLE 8.3. Replacing  $x$  by  $x^2$  in (8.6) gives

$$(8.12) \quad \frac{1 - x^{2n+2}}{(1 - x^2)^2} = \sum_{k=0}^n (k+1)x^{2k} + (n+1) \sum_{k=n+1}^{\infty} x^{2k}.$$

This gives

$$\begin{aligned} \int_0^1 \frac{1 - x^{2n+2}}{(1 - x^2)^2} \ln x \, dx &= \sum_{k=0}^n (k+1) \int_0^1 x^{2k} \ln x \, dx + (n+1) \sum_{k=n+1}^{\infty} \int_0^1 x^{2k} \ln x \, dx \\ &= -\sum_{k=0}^n \frac{k+1}{(2k+1)^2} - (n+1) \sum_{k=n+1}^{\infty} \frac{1}{(2k+1)^2}. \end{aligned}$$

The value

$$(8.13) \quad \sum_{k=1}^{\infty} \frac{1}{(2k-1)^2} = \frac{\pi^2}{8} = \frac{3}{4}\zeta(2)$$

is obtained by separating the terms forming the series for  $\zeta(2)$  into even and odd indices. Now write

$$(8.14) \quad \sum_{k=n+1}^{\infty} \frac{1}{(2k+1)^2} = \frac{3}{4}\zeta(2) - \sum_{k=1}^{n+1} \frac{1}{(2k-1)^2}$$

to obtain, after some elementary algebraic manipulations, the evaluation

$$(8.15) \quad \int_0^1 \frac{1 - x^{2n+2}}{(1 - x^2)^2} \ln x \, dx = -\frac{3}{4}(n+1)\zeta(2) + \sum_{k=1}^n \frac{n-k+1}{(2k-1)^2}.$$

This is entry **4.231.16**.

EXAMPLE 8.4. The alternating geometric series

$$(8.16) \quad \frac{1}{1+x} = \sum_{k=0}^{\infty} (-1)^k x^k$$

is used as before to derive the identity

$$(8.17) \quad \frac{1 + (-1)^n x^{n+1}}{(1+x)^2} = (n+1) \sum_{k=0}^{\infty} (-1)^k x^k - \sum_{k=0}^n (-1)^k (n-k)x^k.$$

Integrating yields

$$(8.18) \quad \int_0^1 \frac{1 + (-1)^n x^{n+1}}{(1+x)^2} \ln x \, dx = -(n+1) \sum_{k=0}^{\infty} \frac{(-1)^k}{(k+1)^2} - \sum_{k=1}^n \frac{(-1)^k (n-k+1)}{k^2}.$$



This is entry **4.231.17**, written in the form

$$(8.19) \quad \int_0^1 \frac{1 + (-1)^n x^{n+1}}{(1+x)^2} \ln x \, dx = -\frac{(n+1)\pi^2}{12} - \sum_{k=1}^n \frac{(-1)^k (n-k+1)}{k^2},$$

using the value

$$(8.20) \quad \sum_{k=0}^{\infty} \frac{(-1)^k}{(k+1)^2} = \frac{\pi^2}{12}.$$

EXAMPLE 8.5. Entry **4.262.8**

$$(8.21) \quad \int_0^1 \frac{1 + (-1)^n x^{n+1}}{(1+x)^2} (\ln x)^3 \, dx = -\frac{7(n+1)\pi^4}{120} + 6 \sum_{k=1}^n (-1)^{k-1} \frac{n-k+1}{k^4}$$

is obtained by using (8.17) and the identities employed in Example 8.2. The procedure employed in Example 8.3 now gives entry **4.262.9**

$$(8.22) \quad \int_0^1 \frac{1 - x^{2n+2}}{(1-x^2)^2} (\ln x)^3 \, dx = -\frac{(n+1)\pi^4}{16} + 6 \sum_{k=1}^n \frac{n-k+1}{(2k-1)^4}.$$

### 9. A singular integral

The last evaluation presented here is entry **4.231.10**

$$(9.1) \quad \int_0^{\infty} \frac{\ln x \, dx}{a^2 - b^2 x^2} = -\frac{\pi^2}{4ab}.$$

The parameters  $a, b$  have the same sign, so it may be assumed that  $a, b > 0$ . Observe that this is a singular integral, since the integrand is discontinuous at  $x = a/b$ .

The change of variables  $t = bx/a$  gives

$$(9.2) \quad \int_0^{\infty} \frac{\ln x \, dx}{a^2 - b^2 x^2} = \frac{1}{ab} \left[ \ln \frac{a}{b} \int_0^{\infty} \frac{dt}{1-t^2} + \int_0^{\infty} \frac{\ln t \, dt}{1-t^2} \right].$$

The first integral is singular and is computed as the limit as  $\varepsilon \rightarrow 0$  of

$$(9.3) \quad \int_0^{1-\varepsilon} \frac{dt}{1-t^2} + \int_{1+\varepsilon}^{\infty} \frac{dt}{1-t^2} = \frac{1}{2} \ln \left( \frac{2-\varepsilon}{\varepsilon} \right) + \frac{1}{2} \ln \left( \frac{\varepsilon}{2+\varepsilon} \right) = \frac{1}{2} \ln \left( \frac{2-\varepsilon}{2+\varepsilon} \right)$$

obtained by the method of partial fraction. Therefore this singular integral has value 0. The second integral is

$$(9.4) \quad \int_0^{\infty} \frac{\ln t \, dt}{1-t^2} = 2 \int_0^1 \frac{\ln t \, dt}{1-t^2},$$

because the integral over  $[1, \infty)$  is the same as over  $[0, 1]$ . The method of partial fractions and the values

$$(9.5) \quad \int_0^1 \frac{\ln x \, dx}{1-x} = -\frac{\pi^2}{6} \quad \text{and} \quad \int_0^1 \frac{\ln x \, dx}{1+x} = -\frac{\pi^2}{12},$$

that appear as entries **4.231.2** and **4.231.1**, respectively, give the final result. These last two entries were evaluated in [1].

The change of variables  $t = \ln x$  converts this integral into entry **3.417.2**

$$(9.6) \quad \int_{-\infty}^{\infty} \frac{t dt}{a^2 e^t - b^2 e^{-t}} = \frac{\pi^2}{4ab}.$$

The same change of variables gives the evaluation of entry **3.417.1**

$$(9.7) \quad \int_{-\infty}^{\infty} \frac{t dt}{a^2 e^t + b^2 e^{-t}} = \frac{\pi}{2ab} \ln \frac{b}{a}$$

from entry **4.231.8**

$$(9.8) \quad \int_0^{\infty} \frac{\ln x dx}{a^2 + b^2 x^2} = -\frac{\pi}{2ab} \ln \frac{b}{a}$$

evaluated in [4].

**Summary.** The examples presented here, complete the evaluation of every entry in Section 4.231 of the table [6]. The entries not appearing here have been presented in [4, 5, 7].

**Acknowledgments.** The second author acknowledges the partial support of NSF-DMS 0713836.

### References

- [1] T. Amdeberhan, K. Boyadzhiev, and V. Moll. The integrals in Gradshteyn and Ryzhik. Part 17: The Riemann zeta function. *Scientia*, 20:61–71, 2011.
- [2] T. Amdeberhan and V. Moll. The integrals in Gradshteyn and Ryzhik. Part 7: Elementary examples. *Scientia*, 16:25–40, 2008.
- [3] T. Amdeberhan and V. Moll. The integrals in Gradshteyn and Ryzhik. Part 14: An elementary evaluation of entry 3.411.5. *Scientia*, 19:97–103, 2010.
- [4] T. Amdeberhan, V. Moll, J. Rosenberg, A. Straub, and P. Whitworth. The integrals in Gradshteyn and Ryzhik. Part 9: Combinations of logarithmic, rational and trigonometric functions. *Scientia*, 17:27–44, 2009.
- [5] K. Boyadzhiev, L. Medina, and V. Moll. The integrals in Gradshteyn and Ryzhik. Part 11: The incomplete beta function. *Scientia*, 18:61–75, 2009.
- [6] I. S. Gradshteyn and M. Ryzhik, I. *Table of Integrals, Series, and Products*. Edited by A. Jeffrey and D. Zwillinger. Academic Press, New York, 7th edition, 2007.
- [7] V. Moll. The integrals in Gradshteyn and Ryzhik. Part 2: Elementary logarithmic integrals. *Scientia*, 14:7–15, 2007.
- [8] V. Moll and R. Posey. The integrals in Gradshteyn and Ryzhik. Part 12: Some logarithmic integrals. *Scientia*, 18:77–84, 2009.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PUERTO RICO, RIO PIEDRAS, PR 00931  
*E-mail address:* [luis.medina17@upr.edu](mailto:luis.medina17@upr.edu)

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LA 70118  
*E-mail address:* [vhm@math.tulane.edu](mailto:vhm@math.tulane.edu)

*Received ??, revised ??*

DEPARTAMENTO DE MATEMÁTICA  
 UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA  
 CASILLA 110-V,  
 VALPARAÍSO, CHILE