

SCIENTIA

Series A: *Mathematical Sciences*, Vol. ?? (2014), ??

Universidad Técnica Federico Santa María

Valparaíso, Chile

ISSN 0716-8446

© Universidad Técnica Federico Santa María 2014

## The integrals in Gradshteyn and Ryzhik. Part 27: More logarithmic examples

Luis A. Medina and Victor H. Moll

ABSTRACT. The table of Gradshteyn and Ryzhik contains many entries where the integrand is a combination of an elementary function and the logarithmic of another function of the same type. This paper presents proofs of some of these. A sample of examples where the elementary function is replaced by an algebraic function is also discussed.

### 1. Introduction

The compendium [5] contains a large collection of evaluation of integrals of the form

$$(1.1) \quad \int_a^b R_1(x) \ln R_2(x) dx$$

where  $R_1$  and  $R_2$  are rational functions. The first paper in this series [9] considered the family

$$(1.2) \quad f_n(a) = \int_0^\infty \frac{\ln^{n-1} x dx}{(x-1)(x+a)}, \text{ for } n \geq 2 \text{ and } a > 0.$$

The function  $f_n(a)$  is given explicitly by

$$(1.3) \quad f_n(a) = \frac{(-1)^n (n-1)!}{1+a} [1 + (-1)^n] \zeta(n) \\ + \frac{1}{n(1+a)} \sum_{j=0}^{\lfloor n/2 \rfloor} \binom{n}{2j} (2^{2j} - 2) (-1)^{j-1} B_{2j} \pi^{2j} (\log a)^{n-2j}.$$

Here  $\zeta(s)$  is the Riemann zeta function and  $B_{2j}$  is the Bernoulli number. In particular, (1.3) shows that  $(1+a)f_n(a)$  is a polynomial in  $\log a$ .

---

2000 *Mathematics Subject Classification*. Primary 33.

*Key words and phrases*. Integrals.

The first author acknowledges the partial support of UPR-FIPI 1890015.00. The second author wishes to acknowledge the partial support of NSF-DMS 0713836.

Other papers in this series [3, 8, 10] and also [6] considered examples of integrals of this type. The results in [3] can be used to provide explicit expressions for an integral of the type considered here, when the poles of the rational function  $R_2$  in (1.1) have real or purely imaginary parts. The present paper is a continuation of this work.

## 2. Some examples involving rational functions

This section considers of integrals of the form

$$(2.1) \quad \int_a^b R_1(x) \ln R_2(x) dx$$

where  $R_1$  and  $R_2$  are rational functions.

**Example 2.1.** Entry 4.234.4 is

$$(2.2) \quad \int_0^\infty \frac{1-x^2}{(1+x^2)^2} \ln x dx = -\frac{\pi}{2}$$

To evaluate this entry, observe that

$$(2.3) \quad \frac{d}{dx} \frac{x}{1+x^2} = \frac{1-x^2}{(1+x^2)^2},$$

and integrating by parts gives

$$(2.4) \quad \int_0^\infty \frac{1-x^2}{(1+x^2)^2} \ln x dx = - \int_0^\infty \frac{dx}{1+x^2} = -\frac{\pi}{2}.$$

**Example 2.2.** Entry 4.234.5 states that

$$(2.5) \quad \int_0^1 \frac{x^2 \ln x dx}{(1-x^2)(1+x^4)} = -\frac{\pi^2}{16(2+\sqrt{2})}.$$

To prove this use the method of partial fraction to obtain

$$(2.6) \quad \int_0^1 \frac{x^2 \ln x dx}{(1-x^2)(1+x^4)} = \frac{1}{4} \int_0^1 \frac{\ln x dx}{1-x} + \frac{1}{4} \int_0^1 \frac{\ln x dx}{1+x} + \frac{1}{2} \int_0^1 \frac{(x^2-1) \ln x dx}{1+x^4}.$$

The first integral is  $-\pi^2/6$  according to entry 4.231.2 and the second one is  $-\pi^2/12$  from entry 4.231.1. These entries were established in [1]. This gives

$$(2.7) \quad \int_0^1 \frac{x^2 \ln x dx}{(1-x^2)(1+x^4)} = -\frac{\pi^2}{16} + \frac{1}{2} \int_0^1 \frac{(x^2-1) \ln x dx}{1+x^4}.$$

To evaluate the last integral, observe that

$$(2.8) \quad \frac{x^2-1}{1+x^4} = \sum_{n=0}^{\infty} (-1)^{n-1} x^{4n} + \sum_{n=0}^{\infty} (-1)^n x^{4n+2}.$$

Now recall the *digamma function*  $\psi(z) = \Gamma'(z)/\Gamma(z)$  and the expansion of its derivative

$$(2.9) \quad \psi'(x) = \sum_{n=0}^{\infty} \frac{1}{(x+n)^2}.$$

Details about this function may be found in [4] and [13]. This gives

$$(2.10) \quad \int_0^1 \frac{(x^2 - 1) \ln x \, dx}{1 + x^4} = \frac{1}{64} \left[ \psi' \left( \frac{1}{8} \right) - \psi' \left( \frac{3}{8} \right) - \psi' \left( \frac{5}{8} \right) + \psi' \left( \frac{7}{8} \right) \right].$$

The classical relation

$$(2.11) \quad \Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin \pi x}$$

can be shifted to produce

$$(2.12) \quad \Gamma\left(\frac{1}{2} + x\right) \Gamma\left(\frac{1}{2} - x\right) = \frac{\pi}{\cos \pi x}.$$

Logarithmic differentiation shows that the digamma function satisfies

$$(2.13) \quad \psi\left(\frac{1}{2} + x\right) - \psi\left(\frac{1}{2} - x\right) = \pi \tan \pi x.$$

This appears as Entry **8.365.9** in [5]. Differentiation produces

$$(2.14) \quad \psi'\left(\frac{1}{2} + x\right) + \psi'\left(\frac{1}{2} - x\right) = \pi^2 \sec^2 \pi x.$$

Now use (2.14) and group 1/8 with 7/8 and 3/8 with 5/8 to produce

$$(2.15) \quad \int_0^1 \frac{(x^2 - 1) \ln x \, dx}{1 + x^4} = \frac{1}{64} \left( \frac{4\pi^2}{2 - \sqrt{2}} - \frac{4\pi^2}{2 + \sqrt{2}} \right) = \frac{\pi^2}{8\sqrt{2}}.$$

**Note 2.3.** The reader should evaluate the family of integrals

$$(2.16) \quad I_n = \int_0^1 \frac{x^{2n} \ln x}{(1 - x^2)(1 + x^4)^n} \, dx, \quad n \in \mathbb{N},$$

by the method described here. The computation of the first few special values indicates an interesting arithmetic structure of the answer.

### 3. An entry involving the Poisson kernel for the disk

The section discusses a single entry in [5], where the integrand involves the Poisson kernel for the disk. Further examples of this type will be presented in a future publication.

**Example 3.1.** The next evaluation is Entry **4.233.5**:

$$(3.1) \quad \int_0^\infty \frac{\ln x \, dx}{x^2 + 2xa \cos t + a^2} = \frac{t \ln a}{\sin t \, a}.$$

The integrand is related to the *Poisson kernel* for the unit disk  $D = \{z \in \mathbb{C} : |z| < 1\}$ .

**Theorem 3.2.** Define

$$(3.2) \quad \mathcal{P}_r(\theta) = \operatorname{Re} \frac{1 + re^{i\theta}}{1 - re^{i\theta}}$$

then  $\mathcal{P}_r(\theta)$  is given by

$$(3.3) \quad \mathcal{P}_r(\theta) = \sum_{n=-\infty}^{\infty} r^{|n|} e^{in\theta} = \frac{1 - r^2}{1 - 2r \cos \theta + r^2}.$$

Moreover, given  $f$  defined on the boundary of  $D$ , the expression

$$(3.4) \quad u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \mathcal{P}_r(\theta - t) f(e^{it}) dt$$

for  $0 \leq r < 1$ , is a harmonic function on  $D$  and it has a radial limit which agrees with  $f$  almost everywhere on the boundary of  $D$ .

The form of the Poisson kernel can be used to establish the next result.

**Lemma 3.3.** For  $a, x \in \mathbb{R}$  with  $|x| < |a|$ ,

$$(3.5) \quad \sum_{k=0}^{\infty} \frac{(-1)^k \sin((k+1)t)x^k}{a^k} = \frac{a^2 \sin t}{x^2 + 2ax \cos t + a^2}.$$

**Note 3.4.** The Chebyshev polynomial of the second kind  $U_n(t)$  is defined by the identity

$$(3.6) \quad \frac{\sin((n+1)\theta)}{\sin \theta} = U_n(\cos \theta).$$

The result of Lemma 3.3 can be written as

$$(3.7) \quad \sum_{k=0}^{\infty} U_k(t)x^k = \frac{1}{x^2 - 2x \cos t + 1}.$$

Lemma 3.3 produces

$$(3.8) \quad \int_0^R \frac{x^s dx}{x^2 + 2ax \cos t + a^2} = \frac{1}{a^2 \sin t} \sum_{k=0}^{\infty} \frac{(-1)^k \sin((k+1)t)R^{k+s+1}}{a^k (k+s+1)}.$$

Now write  $\sin((k+1)t)$  in terms of exponential to obtain an expression for the previous integral as

$$\int_0^R \frac{x^s dx}{x^2 + 2ax \cos t + a^2} = \frac{R^{s+1}}{2ia^2 \sin t} \left( e^{it} \Phi \left( -\frac{R}{ae^{it}}, 1, s+1 \right) - e^{-it} \Phi \left( -\frac{R}{ae^{-it}}, 1, s+1 \right) \right)$$

where

$$(3.9) \quad \Phi(z, s, a) = \sum_{k=0}^{\infty} \frac{z^k}{(a+k)^s}$$

is the *Lerch Phi function*.

Now differentiate with respect to  $s$  and let  $s \rightarrow 0$  to produce

$$(3.10) \quad \int_0^R \frac{\ln x dx}{x^2 + 2ax \cos t + a^2} = \frac{i \ln R}{2a \sin t} (\log(1 + e^{-it}R/a) - \log(1 + e^{it}R/a)) \\ + \frac{i}{2a \sin t} (\text{Li}_2(-e^{-it}R/a) - \text{Li}_2(-e^{it}R/a)),$$

where

$$(3.11) \quad \text{Li}_2(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^2}$$

is the *dilogarithm* function. Then use the identity

$$(3.12) \quad i (\operatorname{Li}_2(-e^{-it}R/a) - \operatorname{Li}_2(-e^{-it}R/a)) = - \int_0^t \ln \left( \frac{a^2 + 2Ra \cos z + R^2}{a^2} \right) dz$$

to obtain

$$(3.13) \quad \int_0^R \frac{\ln x dx}{x^2 + 2ax \cos t + a^2} = \frac{i \ln R}{2a \sin t} (\log(1 + e^{-it}R/a) - \log(1 + e^{it}R/a)) \\ - \frac{1}{2a \sin t} \int_0^t \ln \left( \frac{a^2 + 2Ra \cos z + R^2}{a^2} \right) dz.$$

The next step is to differentiate (3.13) with respect to  $t$  and let  $R \rightarrow \infty$ . The left-hand side produces

$$(3.14) \quad T_1(a, t) = \int_0^\infty \frac{2ax \ln x \sin t dx}{(x^2 + 2ax \cos t + a^2)^2}.$$

Direct differentiation of the right-hand side yields

$$(3.15) \quad T_2(a, t) = \lim_{R \rightarrow \infty} V_1(R; a, t) + V_2(R; a, t)$$

where

$$(3.16) \quad V_1(R; a, t) = \frac{R \ln R (R + a \cos t)}{a \sin t (a^2 + 2aR \cos t + R^2)} - \frac{1}{2a \sin t} \ln \left( \frac{a^2 + 2aR \cos t + R^2}{a^2} \right)$$

and

$$(3.17) \quad V_2(R; a, t) = \frac{i \cos t \ln R}{2a \sin^2 t} (\log(1 + e^{it}R/a) - \log(1 + e^{-it}R/a)) \\ + \frac{\cos t}{2a \sin^2 t} \int_0^t \ln \left( \frac{a^2 + 2Ra \cos z + R^2}{a^2} \right) dz.$$

**Proposition 3.5.** The function  $T_2(a, t)$  is given

$$(3.18) \quad T_2(a, t) = -\frac{\ln a}{2a \sin t} (t \cot t - 1).$$

PROOF. Start with the computation of the limiting behavior of  $V_1(R; a, t)$ . The claim that

$$(3.19) \quad \lim_{R \rightarrow \infty} V_1(R; a, t) = \frac{\ln a}{a \sin(t)}$$

is verified first.

First note that since

$$(3.20) \quad \lim_{R \rightarrow \infty} \frac{R \ln R}{a^2 + 2aR \cos(t) + R^2} = 0,$$

then

$$\lim_{R \rightarrow \infty} V_1(R; a, t) = \frac{1}{a \sin t} \lim_{R \rightarrow \infty} \left( \frac{R^2 \ln R}{a^2 + 2aR \cos t + R^2} - \frac{1}{2} \ln(a^2 + 2aR \cos t + R^2) + \ln a \right).$$

The claim is equivalent to

$$(3.21) \quad \lim_{R \rightarrow \infty} \left( \frac{R^2 \ln R}{a^2 + 2aR \cos t + R^2} - \frac{1}{2} \ln(a^2 + 2aR \cos t + R^2) \right) = 0.$$

The identities

$$(3.22) \quad \frac{R^2 \ln R}{a^2 + 2aR \cos t + R^2} = \frac{\ln R}{a^2/R^2 + 2a \cos t/R + 1}$$

and

$$(3.23) \quad \frac{1}{2} \ln(a^2 + 2aR \cos t + R^2) = \ln R + \frac{1}{2} \ln(a^2/R^2 + 2a \cos t/R + 1)$$

can be used to see that the left-hand side of (3.21) is equivalent to

$$\lim_{R \rightarrow \infty} \left( \ln R \left( \frac{1}{a^2/R^2 + 2a \cos t/R + 1} - 1 \right) - \frac{1}{2} \ln(a^2/R^2 + 2a \cos t/R + 1) \right) = 0.$$

It is clear that the second term vanishes as  $R \rightarrow \infty$ . For the first term, observe that

$$(3.24) \quad \frac{1}{a^2/R^2 + 2a \cos(t)/R + 1} - 1 = -\frac{2a \cos t}{R} + O\left(\frac{1}{R^2}\right)$$

and thus the first term also vanishes as  $R \rightarrow \infty$ . This concludes the proof.

The next step is to verify that

$$(3.25) \quad V_2(R; a, t) = \frac{i \cot t \ln R}{2a \sin^2 t} (\log(1 + e^{it} R/a) - \log(1 + e^{-it} R/a)) \\ + \frac{\cos t}{2a \sin^2 t} \int_0^t \ln \left( \frac{a^2 + 2aR \cos z + R^2}{a^2} \right) dz$$

satisfies

$$(3.26) \quad \lim_{R \rightarrow \infty} V_2(R; a, t) = -\frac{t \cos t}{a \sin^2 t} \ln a.$$

The proof begins with the identity

$$(3.27) \quad \log(1 + b/x) = \log(b/x) + \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{nb^n}$$

to obtain

$$(3.28) \quad \log(1 + e^{it} R/a) - \log(1 + e^{-it} R/a) = \log(e^{it}) - \log(e^{-it}) + O(a/R), \text{ as } R \rightarrow \infty.$$

The bounds  $0 < t < \pi$  imply  $\log(e^{it}) - \log(e^{-it}) = 2it$ . This gives

$$\begin{aligned} \lim_{R \rightarrow \infty} V_2(R; a, t) &= \lim_{R \rightarrow \infty} \left( \frac{\cos t}{2a \sin^2 t} \int_0^t \ln \left( \frac{a^2 + 2aR \cos z + R^2}{a^2} \right) dz - \frac{t \cos z \ln R}{a \sin^2 t} \right) \\ &= \lim_{R \rightarrow \infty} \frac{\cos t}{2a \sin^2 t} \left( \int_0^t \ln \left( \frac{a^2 + 2aR \cos z + R^2}{a^2} \right) dz - 2t \ln R \right) \\ &= \lim_{R \rightarrow \infty} \frac{\cos t}{2a \sin^2 t} \left( \int_0^t \ln \left( \frac{a^2 + 2aR \cos z + R^2}{a^2} \right) - \ln(R^2) dz \right) \\ &= \lim_{R \rightarrow \infty} \frac{\cos t}{2a \sin^2 t} \left( \int_0^t [\ln(a^2 + 2aR \cos z + R^2) - \ln(R^2)] dz - 2t \ln a \right). \end{aligned}$$

The identity

$$\int_0^t [\ln(a^2 + 2aR \cos z + R^2) - \ln(R^2)] dz = \int_0^t \ln \left( \frac{a^2}{R^2} + \frac{2a \cos z}{R} + 1 \right) dz$$

gives the result. The proof of the Proposition is finished.  $\square$

The evaluation of entry **4.233.5** is now obtained from the identity  $T_1(a, t) = T_2(a, t)$ . Observe that this implies

$$(3.29) \quad \int_0^\infty \frac{2ax \ln x \sin t dx}{(x^2 + 2ax \cos t + a^2)^2} = -\frac{\ln a}{a \sin t} (t \cot t - 1).$$

Integrating with respect to  $t$  gives (3.1). Entry **4.231.8** in [5], established in [3],

$$(3.30) \quad \int_0^\infty \frac{\ln x dx}{x^2 + a^2} = \frac{\pi \ln a}{2a}$$

can be used to show that the implicit constant of integration actually vanishes. The evaluation is complete.

#### 4. Some rational integrands with a pole at $x = 1$

This section contains proofs of the four entries appearing in Section 4.235. These are integrals of the form

$$(4.1) \quad f(a, b, c) := \int_0^\infty \frac{x^b - x^c}{1 - x^a} \ln x dx$$

where  $a, b, c \in \mathbb{N}$ . These integrals are evaluated using entry **4.254.2**

$$(4.2) \quad \int_0^\infty \frac{x^{p-1} \ln x}{1 - x^q} dx = -\frac{\pi^2}{q^2 \sin^2 \frac{\pi p}{q}}.$$

To obtain this formula, start from **3.231.6**

$$(4.3) \quad \int_0^\infty \frac{x^{p-1} - x^{q-1}}{1 - x} dx = \pi (\cot \pi p - \cot \pi q),$$

established in [7] and make the change of variables  $t = x^q$  to produce

$$\begin{aligned} \int_0^\infty \frac{x^{p-1} - 1}{1 - x^q} dx &= -\frac{1}{q} \int_0^\infty \frac{t^{1/q-1} - t^{p/q-1}}{1 - t} dt \\ &= -\frac{\pi}{q} \left( \cot \frac{\pi}{q} - \cot \frac{\pi p}{q} \right). \end{aligned}$$

Differentiating with respect to  $p$  gives (4.2).

LEMMA 4.1. *Let  $a, b, c \in \mathbb{R}$ . Then*

$$(4.4) \quad \int_0^\infty \frac{x^{b-1} - x^{c-1}}{1 - x^a} \ln x dx = -\frac{\pi^2}{a^2} \frac{\sin(c_1 - b_1) \sin(c_1 + b_1)}{\sin^2 b_1 \sin^2 c_1}$$

where  $b_1 = \pi b/a$  and  $c_1 = \pi c/a$ .

PROOF. Simply write

$$\int_0^\infty \frac{x^{b-1} - x^{c-1}}{1 - x^a} \ln x dx = \int_0^\infty \frac{x^{b-1}}{1 - x^a} \ln x dx - \int_0^\infty \frac{x^{c-1}}{1 - x^a} \ln x dx$$

and use (4.2). □

The four entries in Section 4.235 are established next.

**Example 4.1.** Entry 4.235.1 states that

$$(4.5) \quad \int_0^\infty \frac{(1-x)x^{n-2}}{1-x^{2n}} \ln x dx = -\frac{\pi^2}{4n^2} \tan^2 \frac{\pi}{2n}.$$

Lemma 4.1 is used with  $a = 2n$ ,  $b = n - 1$  and  $c = n$ . This gives

$$(4.6) \quad b_1 = \frac{\pi}{2} - \frac{\pi}{2n} \text{ and } c_1 = \frac{\pi}{2}.$$

and

$$\int_0^\infty \frac{(1-x)x^{n-2}}{1-x^{2n}} \ln x dx = -\frac{\pi^2}{4n^2} \frac{\sin\left(\frac{\pi}{2} - \frac{\pi}{2n}\right) \sin\left(\frac{\pi}{2} + \frac{\pi}{2n}\right)}{\sin^2\left(\frac{\pi}{2} - \frac{\pi}{2n}\right)} = -\frac{\pi^2}{4n^2} \tan^2 \frac{\pi}{2n}.$$

**Example 4.2.** Entry 4.235.2 is

$$(4.7) \quad \int_0^\infty \frac{(1-x^2)x^{m-1}}{1-x^{2n}} \ln x dx = -\frac{\pi^2}{4n^2} \frac{\sin\left(\frac{m+1}{n}\pi\right) \sin\left(\frac{\pi}{n}\right)}{\sin^2\left(\frac{\pi m}{2n}\right) \sin^2\left(\frac{(m+2)}{2n}\pi\right)}.$$

Lemma 4.1 is now used with  $a = 2n$ ,  $b = m$  and  $c = m + 2$ . This gives

$$(4.8) \quad c_1 - b_1 = \frac{\pi}{n} \text{ and } c_1 + b_1 = \frac{\pi}{n}(m + 1)$$

to produce the result.

**Example 4.3.** Entry 4.235.3 states that

$$(4.9) \quad \int_0^\infty \frac{(1-x^2)x^{n-3}}{1-x^{2n}} \ln x dx = -\frac{\pi^2}{4n^2} \tan^2 \frac{\pi}{n}.$$

The values  $a = 2n$ ,  $b = n - 2$  and  $c = n$  give

$$(4.10) \quad b_1 = \frac{\pi}{2} - \frac{\pi}{n} \text{ and } c_1 = \frac{\pi}{2}.$$



This verifies the claim.

**Example 4.4.** Entry 4.235.4 appears as

$$(4.11) \quad \int_0^1 \frac{x^{m-1} + x^{n-m-1}}{1-x^n} \ln x \, dx = -\frac{\pi^2}{n^2 \sin^2 \frac{\pi m}{n}}.$$

The change of variables  $t = 1/x$  shows that the integral over  $[1, \infty)$  is equal to that over  $[0, 1]$ , therefore this entry should be written as

$$(4.12) \quad \int_0^\infty \frac{x^{m-1} + x^{n-m-1}}{1-x^n} \ln x \, dx = -\frac{2\pi^2}{n^2 \sin^2 \frac{\pi m}{n}},$$

to be consistent with the other entries in this section. The proof comes from Lemma 4.1 with  $a = n$ ,  $b = m$  and  $c = n - m$ .

### 5. Some singular integrals

The table [5] contains a variety of singular integrals of the form being discussed here. The examples considered in this section are evaluated employing the formula

$$(5.1) \quad \int_0^\infty \frac{t^{\mu-1} dt}{1-t} = \pi \cot \pi \mu.$$

To verify this evaluation, transform the integral over  $[1, \infty)$  to  $[0, 1]$  by the change of variables  $x \mapsto 1/x$ . This gives

$$(5.2) \quad \int_0^\infty \frac{t^{\mu-1} dt}{1-t} = \int_0^1 \frac{t^{\mu-1} - t^{-\mu}}{1-t} dt.$$

This is entry 3.231.1. It was established in [7].

Differentiating with respect to  $\mu$ , the formula (5.1) gives

$$(5.3) \quad \int_0^\infty \frac{t^{\mu-1} \ln t}{1-t} dt = -\frac{\pi^2}{\sin^2 \pi \mu},$$

and the change of variables  $t = x^a$  gives

$$(5.4) \quad \omega(a, b) := \int_0^\infty \frac{x^{b-1} \ln x}{1-x^a} dx = -\frac{\pi^2}{a^2 \sin^2 \left(\frac{\pi b}{a}\right)}.$$

**Example 5.1.** Entry 4.251.2 states that

$$(5.5) \quad \int_0^\infty \frac{x^{\mu-1} \ln x}{a-x} = \pi a^{\mu-1} \left( \ln a \cot(\pi \mu) - \frac{\pi}{\sin^2 \pi \mu} \right).$$

The change of variables  $x = at$  yields

$$(5.6) \quad \int_0^\infty \frac{x^{\mu-1} \ln x}{a-x} = a^{\mu-1} \int_0^\infty \frac{t^{\mu-1} \ln t}{1-t} dt + a^{\mu-1} \ln a \int_0^\infty \frac{t^{\mu-1} dt}{1-t}.$$

The result now follows from (5.1) and (5.3). It is probably clearer to write this entry as

$$(5.7) \quad \int_0^\infty \frac{x^{\mu-1} \ln x}{a-x} = \pi a^{\mu-1} \left( \frac{\ln a}{\tan \pi \mu} - \frac{\pi}{\sin^2 \pi \mu} \right),$$

to avoid possible confusions.

**Example 5.2.** Entry 4.252.3 is

$$(5.8) \quad \int_0^\infty \frac{x^{p-1} \ln x}{1-x^2} dx = -\frac{\pi^2}{4} \operatorname{cosec}^2 \frac{\pi p}{2}.$$

This is  $\omega(2, p)$  and the result follows from (5.4).

**Example 5.3.** Entry 4.255.3 states that

$$(5.9) \quad \int_0^\infty \frac{1-x^p}{1-x^2} \ln x dx = \frac{\pi^2}{4} \tan^2 \left( \frac{\pi p}{2} \right).$$

This is  $\omega(1, 2) - \omega(p+1, 2)$  and the result comes from (5.4).

**Example 5.4.** Entry 4.252.1 is written as

$$\int_0^\infty \frac{x^{\mu-1} \ln x dx}{(x+a)(x+b)} = \frac{\pi}{(b-a) \sin \pi \mu} \left[ a^{\mu-1} \ln a - b^{\mu-1} \ln b - \pi \frac{a^{\mu-1} - b^{\mu-1}}{\tan \pi \mu} \right].$$

This value follows from the partial fraction decomposition

$$(5.10) \quad \frac{1}{(x+a)(x+b)} = \frac{1}{b-a} \frac{1}{x+a} - \frac{1}{b-a} \frac{1}{x+b}$$

and entry 4.251.1

$$(5.11) \quad \int_0^\infty \frac{x^{\mu-1} \ln x}{x+c} dx = \frac{\pi c^{\mu-1}}{\sin \pi \mu} (\ln c - \pi \cot \pi \mu),$$

established in [11]. Differentiating (5.11) with respect to  $c$  yields

$$(5.12) \quad \int_0^\infty \frac{x^{\mu-1} \ln x}{(x+c)^2} dx = -\frac{(\mu-1)c^{\mu-2}\pi}{\sin \pi \mu} \left( \ln c - \pi \cot \pi \mu + \frac{1}{\mu-1} \right).$$

This is entry 4.252.4.

**Example 5.5.** Entry 4.257.1

$$(5.13) \quad \int_0^\infty \frac{x^\mu \ln(x/a) dx}{(x+a)(x+b)} = \frac{\pi [b^\mu \ln(b/a) + \pi(a^\mu - b^\mu) \cot \pi \mu]}{(b-a) \sin \pi \mu}$$

follows from (5.11) and the beta integral

$$(5.14) \quad \int_0^\infty \frac{x^{\mu-1} dx}{x+a} = \frac{\pi a^{\mu-1}}{\sin \pi \mu}.$$

This appears as entry 3.194.3 and it was established in [11].

**Example 5.6.** The change of variables  $t = x^q$  gives

$$(5.15) \quad \int_0^\infty \frac{x^{p-1} dx}{1-x^q} = \frac{1}{q} \int_0^\infty \frac{t^{p/q-1} dx}{1-t} = \frac{\pi}{q} \cot \left( \frac{\pi p}{q} \right)$$

from (5.3). This is entry 3.241.3. The special case  $q = 1$  gives

$$(5.16) \quad \int_0^\infty \frac{x^{p-1} dx}{1-x} = \pi \cot \pi p.$$

Differentiating with respect to  $p$  produces

$$(5.17) \quad \int_0^\infty \frac{x^{p-1} \ln x}{1-x} dx = -\frac{\pi^2}{\sin^2 \pi p}.$$

The partial fraction decomposition

$$(5.18) \quad \frac{1}{(x+a)(x-1)} = \frac{1}{a+1} \frac{1}{x-1} - \frac{1}{a+1} \frac{1}{x+a}$$

then produces entry **4.252.2**

$$(5.19) \quad \int_0^\infty \frac{x^{\mu-1} \ln x}{(x+a)(x-1)} dx = \frac{\pi}{(a+1) \sin^2 \pi \mu} [\pi - a^{\mu-1} (\ln a \sin \pi \mu - \pi \cos \pi \mu)].$$

**Example 5.7.** The change of variables  $t = x^q$  produces

$$(5.20) \quad \int_0^\infty \frac{\ln x dx}{x^p(x^q-1)} = -\frac{1}{q^2} \int_0^\infty \frac{t^{(1-p)/q-1} \ln t dt}{1-t}.$$

Then, (5.3) gives

$$(5.21) \quad \int_0^\infty \frac{\ln x dx}{x^p(x^q-1)} = \frac{\pi^2}{q^2} \frac{1}{\sin^2 \left( \frac{p-1}{q} \pi \right)}.$$

This is entry **4.254.3**.

**Example 5.8.** Entry **4.255.2** is

$$(5.22) \quad \int_0^1 \frac{(1+x^2)x^{p-2}}{1-x^{2p}} \ln x dx = -\left(\frac{\pi}{2p}\right)^2 \sec^2 \frac{\pi}{2p}.$$

The evaluation of this entry starts with entry **3.231.5**

$$(5.23) \quad \int_0^1 \frac{x^{\mu-1} - x^{\nu-1}}{1-x} dx = -\psi(\mu) + \psi(\nu)$$

that was established in [7]. The special case  $\mu = 1$

$$(5.24) \quad \int_0^1 \frac{1-x^{\nu-1}}{1-x} dx = -\psi(1) + \psi(\nu)$$

is differentiated with respect to  $\nu$  to produce

$$(5.25) \quad \int_0^1 \frac{x^{\nu-1} \ln x}{1-x} dx = -\psi'(\nu).$$

The change of variables  $x = t^b$  gives

$$(5.26) \quad \int_0^1 \frac{t^{c-1} \ln t}{1-t^b} dt = -\frac{1}{b^2} \psi' \left( \frac{c}{b} \right).$$

Therefore

$$\begin{aligned} \int_0^1 \frac{(1-x^2)x^{p-2}}{1-x^{2p}} \ln x dx &= \int_0^1 \frac{x^{p-2}}{1-x^{2p}} \ln x dx + \int_0^1 \frac{x^p}{1-x^{2p}} \ln x dx \\ &= -\frac{1}{4p^2} \left[ \psi' \left( \frac{1}{2} - \frac{1}{2p} \right) + \psi' \left( \frac{1}{2} + \frac{1}{2p} \right) \right]. \end{aligned}$$

The result now follows from the reflection formula for the polygamma function  $\psi'$  given in (2.14).

### 6. Combinations of logarithms and algebraic functions

This section presents the evaluation of some entries in [5] of the form

$$(6.1) \quad \int_a^b E_1(x) \ln E_2(x) dx$$

where  $E_1$  or  $E_2$  is an algebraic function. Some of these have appeared in previous papers in this series. For example, entry **4.241.11**

$$(6.2) \quad \int_0^1 \frac{\ln x dx}{\sqrt{x(1-x^2)}} = -\frac{\sqrt{2\pi}}{8} \Gamma^2\left(\frac{1}{4}\right)$$

and entry **4.241.5**

$$(6.3) \quad \int_0^1 \ln x \sqrt{(1-x^2)^{2n-1}} dx = -\frac{(2n-1)!!}{4(2n)!!} \pi [\psi(n+1) + \gamma + \ln 4]$$

were evaluated in [7]. Here  $\psi(x)$  is the digamma function and  $\gamma$  is Euler's constant.

**Note 6.1.** Define the family of integrals

$$(6.4) \quad f_n(a) := \int_0^1 \frac{x^a \ln^n x dx}{\sqrt{1-x^2}}.$$

Special cases include entry **4.241.7**

$$(6.5) \quad \int_0^1 \frac{\ln x dx}{\sqrt{1-x^2}} = -\frac{\pi}{2} \ln 2$$

that was evaluated in [7] and entry **4.261.9**

$$(6.6) \quad \int_0^1 \frac{\ln^2 x dx}{\sqrt{1-x^2}} = \frac{\pi}{2} \left( \ln^2 2 + \frac{\pi^2}{12} \right).$$

A trigonometric form of the family is obtained by the change of variables  $x = \sin t$ :

$$(6.7) \quad f_n(a) = \int_0^{\pi/2} \sin^a t \ln^n \sin t dt.$$

**Theorem 6.2.** The integral  $f_n(a)$  is given by

$$(6.8) \quad f_n(a) = \lim_{s \rightarrow a} \left( \frac{d}{ds} \right)^n h(s),$$

where

$$(6.9) \quad h(s) = \int_0^{\pi/2} \sin^s t dt = \frac{1}{2} B\left(\frac{s+1}{2}, \frac{1}{2}\right) = \frac{\sqrt{\pi}}{2} \frac{\Gamma\left(\frac{s}{2} + \frac{1}{2}\right)}{\Gamma\left(\frac{s}{2} + 1\right)}.$$

This appears as entry **3.621.5**. Therefore, the evaluation of  $f_n(a)$  requires the values of  $\Gamma^{(k)}(x)$  for  $0 \leq k \leq n$  at  $x = (a+1)/2$  and  $x = a/2 + 1$ .

**Example 6.3.** For example,

$$\begin{aligned} f_1(0) &= \int_0^1 \frac{\ln x \, dx}{\sqrt{1-x^2}} = \lim_{s \rightarrow 0} \frac{d}{ds} \left[ \frac{\sqrt{\pi} \Gamma\left(\frac{s}{2} + \frac{1}{2}\right)}{2 \Gamma\left(\frac{s}{2} + 1\right)} \right] \\ &= \frac{\sqrt{\pi} \Gamma'(1/2)\Gamma(1) - \Gamma'(1)\Gamma(1/2)}{4 \Gamma^2(1)}. \end{aligned}$$

The values

$$(6.10) \quad \Gamma'\left(\frac{1}{2}\right) = -\sqrt{\pi}(\gamma + 2 \ln 2), \Gamma'(1) = -\gamma, \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi} \text{ and } \Gamma(1) = 1$$

give

$$(6.11) \quad f_1(0) = -\frac{\pi}{2} \ln 2.$$

**Proposition 6.4.** The derivatives of the gamma function satisfy the recurrence

$$(6.12) \quad \Gamma^{(n+1)}(x) = \sum_{k=0}^n \binom{n}{k} \Gamma^{(k)}(x) \psi^{(n-k)}(x).$$

**Example 6.5.** A direct application of formula (6.8) evaluates entry **4.261.9**

$$(6.13) \quad f_2(0) = \int_0^1 \frac{\ln^2 x \, dx}{\sqrt{1-x^2}}.$$

Indeed, using  $\Gamma(1) = 1$ , gives

$$(6.14) \quad f_2(0) = \frac{\sqrt{\pi}}{2} \left[ -\frac{1}{2} \Gamma'\left(\frac{1}{2}\right) \Gamma'(1) + \frac{1}{2} \Gamma\left(\frac{1}{2}\right) \Gamma'(1)^2 + \frac{1}{4} \Gamma''\left(\frac{1}{2}\right) - \frac{1}{4} \Gamma\left(\frac{1}{2}\right) \Gamma''(1) \right].$$

The values

$$(6.15) \quad \Gamma''(1) = \gamma^2 + \frac{\pi^2}{6} \text{ and } \Gamma''\left(\frac{1}{2}\right) = \frac{1}{2} \pi^{5/2} + \sqrt{\pi}(\gamma + 2 \ln 2)^2$$

give the identity (6.6).

It remains to explain the values given in (6.10) and (6.15). The recurrence (6.12) reduces the computation of the derivatives of  $\Gamma(x)$  to those of  $\psi(x)$ . The special values given above come from the next result.

**Lemma 6.6.** The digamma function satisfies

$$\begin{aligned} \psi^{(n)}(1) &= (-1)^{n+1} n! \zeta(n+1) \\ \psi^{(n)}\left(\frac{1}{2}\right) &= (-1)^{n+1} n! (2^{n+1} - 1) \zeta(n+1). \end{aligned}$$

PROOF. This comes directly from (2.9). □

**Example 6.7.** The values given in Lemma 6.6 yield

$$\begin{aligned} f_3(0) &= \int_0^1 \frac{\ln^3 x \, dx}{\sqrt{1-x^2}} = -\frac{\pi}{8} (\pi^2 \ln 2 + 4 \ln^3 2 + 6 \zeta(3)) \\ f_4(0) &= \int_0^1 \frac{\ln^4 x \, dx}{\sqrt{1-x^2}} = \frac{\pi}{480} (19\pi^4 + 120\pi^2 \ln^2 2 + 240 \ln^4 2 + 1440 \ln 2 \zeta(3)) \end{aligned}$$

and

$$\begin{aligned} f_1\left(\frac{1}{2}\right) &= \int_0^1 \frac{\sqrt{x} \ln x \, dx}{\sqrt{1-x^2}} = \frac{(\pi-4)}{\sqrt{2\pi}} \Gamma^2\left(\frac{3}{4}\right) \\ f_2\left(\frac{1}{2}\right) &= \int_0^1 \frac{\sqrt{x} \ln^2 x \, dx}{\sqrt{1-x^2}} = \frac{1}{2\sqrt{2\pi}} \Gamma^2\left(\frac{3}{4}\right) (32 - 16G + \pi(\pi-8)), \end{aligned}$$

where  $G$  is **Catalan's constant**

$$(6.16) \quad G = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2}.$$

**Example 6.8.** Entry 4.261.15 states that

$$(6.17) \quad \int_0^1 \frac{x^{2n} \ln^2 x \, dx}{\sqrt{1-x^2}} = \frac{(2n-1)!!}{2(2n)!!} \pi \left\{ \frac{\pi^2}{12} + \sum_{k=1}^{2n} \frac{(-1)^k}{k^2} + \left[ \sum_{k=1}^{2n} \frac{(-1)^k}{k} + \ln 2 \right]^2 \right\}.$$

This is obtained by differentiating  $h(s)$  twice with respect to  $s$  to produce

$$\begin{aligned} \int_0^1 \frac{x^s \ln^2 x \, dx}{\sqrt{1-x^2}} &= \\ &= \frac{\sqrt{\pi}}{8} \frac{\Gamma\left(\frac{s+1}{2}\right)}{\Gamma\left(\frac{s}{2}+1\right)} \left[ \left( \psi\left(\frac{s}{2}+1\right) - \psi\left(\frac{s+1}{2}\right) \right)^2 + \psi'\left(\frac{s+1}{2}\right) - \psi'\left(\frac{s}{2}+1\right) \right]. \end{aligned}$$

Therefore

$$\int_0^1 \frac{x^{2n} \ln^2 x \, dx}{\sqrt{1-x^2}} = \frac{\sqrt{\pi}}{8} \frac{\Gamma\left(n+\frac{1}{2}\right)}{\Gamma(n+1)} \left[ \left( \psi(n+1) - \psi\left(n+\frac{1}{2}\right) \right)^2 + \psi'\left(n+\frac{1}{2}\right) - \psi'(n+1) \right].$$

The special values

$$(6.18) \quad \Gamma\left(n+\frac{1}{2}\right) = \frac{(2n-1)!!}{2^n} \sqrt{\pi} \text{ and } \Gamma(n+1) = n!$$

give

$$\int_0^1 \frac{x^{2n} \ln^2 x \, dx}{\sqrt{1-x^2}} = \frac{\pi}{8} \frac{(2n-1)!!}{(2n)!!} \left[ \left( \psi(n+1) - \psi\left(n+\frac{1}{2}\right) \right)^2 + \psi'\left(n+\frac{1}{2}\right) - \psi'(n+1) \right].$$

Now use the special values

$$(6.19) \quad \psi(n+1) = -\gamma + \sum_{k=1}^n \frac{1}{k} \text{ and } \psi\left(n+\frac{1}{2}\right) = -\gamma - 2\ln 2 + 2 \sum_{k=1}^n \frac{1}{2k-1}$$

as well as

$$(6.20) \quad \psi'(n+1) = \frac{\pi^2}{6} - \sum_{k=1}^n \frac{1}{k^2} \text{ and } \psi'\left(n+\frac{1}{2}\right) = \frac{\pi^2}{2} - 4 \sum_{k=1}^n \frac{1}{(2k-1)^2}$$

to obtain

$$(6.21) \quad \psi(n+1) - \psi\left(n + \frac{1}{2}\right) = 2 \sum_{k=1}^{2n} \frac{(-1)^k}{k} + 2 \ln 2$$

and

$$(6.22) \quad \psi'(n + \frac{1}{2}) - \psi'(n+1) = \frac{\pi^2}{3} + 4 \sum_{k=1}^{2n} \frac{(-1)^k}{k^2}.$$

This gives the result.

**Example 6.9.** A similar analysis gives entry **4.261.16**

$$\int_0^1 \frac{x^{2n+1} \ln^2 x}{\sqrt{1-x^2}} dx = -\frac{(2n)!!}{(2n+1)!!} \left\{ \frac{\pi^2}{12} + \sum_{k=1}^{2n+1} \frac{(-1)^k}{k^2} - \left[ \sum_{k=1}^{2n+1} \frac{(-1)^k}{k} + \ln 2 \right]^2 \right\}.$$

**Example 6.10.** Entry **4.241.6** states that

$$(6.23) \quad \int_0^{1/\sqrt{2}} \frac{\ln x dx}{\sqrt{1-x^2}} = -\frac{\pi}{4} \ln 2 - \frac{G}{2}.$$

The change of variables  $x = \sin t$  gives

$$(6.24) \quad \int_0^{1/\sqrt{2}} \frac{\ln x dx}{\sqrt{1-x^2}} = \int_0^{\pi/4} \ln \sin t dt.$$

This integral is entry **4.224.2** and it has been evaluated in [3].

### 7. An example producing a trigonometric answer

The next example contains, in the logarithmic part, a quotient of linear functions. The evaluation of this entry requires a different approach.

**Example 7.1.** Entry **4.297.8** states that

$$(7.1) \quad \int_0^1 \ln \frac{1+ax}{1-ax} \frac{dx}{x\sqrt{1-x^2}} = \pi \sin^{-1} a.$$

This evaluation starts with the expansion

$$(7.2) \quad \frac{1}{x} \ln \frac{1+ax}{1-ax} = \sum_{n=0}^{\infty} \frac{2a^{2n+1}}{2n+1} x^{2n}$$

to obtain

$$(7.3) \quad \int_0^1 \ln \frac{1+ax}{1-ax} \frac{dx}{x\sqrt{1-x^2}} = \sum_{n=0}^{\infty} \frac{2a^{2n+1}}{2n+1} \int_0^1 \frac{x^{2n} dx}{\sqrt{1-x^2}}.$$

The change of variables  $x = \sin \theta$  gives

$$(7.4) \quad \int_0^1 \frac{x^{2n} dx}{\sqrt{1-x^2}} = \int_0^{\pi/2} \sin^{2n} \theta d\theta = \frac{\pi}{2^{2n+1}} \binom{2n}{n}.$$

The last evaluation is the famous Wallis' formula. It appears as entry **3.621.3** and it was established in [2] and [12]. Therefore

$$(7.5) \quad \int_0^1 \ln \frac{1+ax}{1-ax} \frac{dx}{x\sqrt{1-x^2}} = \sum_{n=0}^{\infty} \frac{\pi}{2^{2n}} \frac{a^{2n+1}}{2n+1} \binom{2n}{n}.$$

The series is now identified from the classical expansion

$$\begin{aligned} \sin^{-1} x &= \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n}{(2n+1)n!} x^{2n+1} \\ &= \sum_{n=0}^{\infty} \frac{1}{2^{2n}(2n+1)} \binom{2n}{n} x^{2n+1} \end{aligned}$$

obtained by expanding the integrand in

$$(7.6) \quad \sin^{-1} x = \int_0^x \frac{dt}{\sqrt{1-t^2}}.$$

as a binomial series and integrating term by term.

Further examples in [5], of the class considered here, will be presented in a future publication.

**Acknowledgments.** The second author acknowledges the partial support of NSF-DMS 0713836.

## References

- [1] T. Amdeberhan, K. Boyadzhiev, and V. Moll. The integrals in Gradshteyn and Ryzhik. Part 17: The Riemann zeta function. *Scientia*, 20:61–71, 2011.
- [2] T. Amdeberhan, L. A. Medina, and V. Moll. The integrals in Gradshteyn and Ryzhik. Part 5: Some trigonometric integrals. *Scientia*, 15:47–60, 2007.
- [3] T. Amdeberhan, V. Moll, J. Rosenberg, A. Straub, and P. Whitworth. The integrals in Gradshteyn and Ryzhik. Part 9: Combinations of logarithmic, rational and trigonometric functions. *Scientia*, 17:27–44, 2009.
- [4] G. Boros and V. Moll. *Irresistible Integrals*. Cambridge University Press, New York, 1st edition, 2004.
- [5] I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products*. Edited by A. Jeffrey and D. Zwillinger. Academic Press, New York, 7th edition, 2007.
- [6] L. Medina and V. Moll. A class of logarithmic integrals. *Ramanujan Journal*, 20:91–126, 2009.
- [7] L. Medina and V. Moll. The integrals in Gradshteyn and Ryzhik. Part 10: The digamma function. *Scientia*, 17:45–66, 2009.
- [8] L. Medina and V. Moll. The integrals in Gradshteyn and Ryzhik. Part 23: Combinations of logarithms and rational functions. *Scientia*, 23:1–18, 2012.
- [9] V. Moll. The integrals in Gradshteyn and Ryzhik. Part 1: A family of logarithmic integrals. *Scientia*, 14:1–6, 2007.



- [10] V. Moll. The integrals in Gradshteyn and Ryzhik. Part 2: Elementary logarithmic integrals. *Scientia*, 14:7–15, 2007.
- [11] V. Moll. The integrals in Gradshteyn and Ryzhik. Part 6: The beta function. *Scientia*, 16:9–24, 2008.
- [12] V. Moll. The integrals in Gradshteyn and Ryzhik. Part 13: Trigonometric forms of the beta function. *Scientia*, 19:91–96, 2010.
- [13] N. M. Temme. *Special Functions. An introduction to the Classical Functions of Mathematical Physics*. John Wiley and Sons, New York, 1996.

DEPARTAMENT OF MATHEMATICS, UNIVERSITY OF PUERTO RICO, RIO PIEDRAS, SAN JUAN, PR 00931

*E-mail address:* `luis.medina17@upr.edu`

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LA 70118

*E-mail address:* `vhm@math.tulane.edu`

*Received ??, revised ??*

DEPARTAMENTO DE MATEMÁTICA  
UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA  
CASILLA 110-V,  
VALPARAÍSO, CHILE